Community Plan of Rhode Island

Quick reference guide

Opioid prevention and opioid use disorder treatment programs for UnitedHealthcare Community Plan of Rhode Island

In response to the U.S. opioid epidemic, we've developed programs to help our members receive the care and treatment they need in safe and effective ways. We've based our measures on the Centers for Disease Control and Prevention's (CDC) opioid treatment guidelines to help prevent overuse of short-acting and long-acting opioid medications. Please use this quick reference guide for information on what we offer.

Concurrent Drug Utilization Review (cDUR) programs

The cDUR program uses the pharmacy claims processing system to screen all prescriptions at the point of service and checks for possible inappropriate drug prescribing and utilization, as well as potentially dangerous medical implications or drug interactions. The program includes communication to the dispensing pharmacy at point of service through claims edits and messaging to the dispensing pharmacy at point of service. The pharmacist will need to address the clinical situation at the point of sale before entering appropriate NCPDP codes to receive an approved claim, unless otherwise stated below.

THERDOSE Acetaminophen	 Combination opioids plus acetaminophen (APAP) limit Prevents doses of APAP greater than 4 grams per day 	
Duplicate Therapy – Short-Acting Opioids (SAOs)	Alerts to concurrent use of multiple SAOs	
Duplicate Therapy – Long-Acting Opioids (LAOs)	Alerts to concurrent use of multiple LAOs	
Drug-Drug Interaction – Opioids and Medication-Assisted Treatment (MAT)	Point-of-sale alert for concurrent use of opioids and MAT drugs	
Drug-Drug Interaction – Opioids and Benzodiazepines	Point-of-sale alert for concurrent use of opioids and benzodiazepines	
Drug-Drug Interaction – Opioids and Carisoprodol	Point-of-sale alert for concurrent use of opioids and carisoprodol	
Drug-Drug Interaction – Opioids and Antipsychotics	Point-of-sale alert for concurrent use of opioids and antipsychotics	
Drug-Drug Interaction – Opioids and Sedative Hypnotics	Point-of-sale alert for concurrent use of opioids and sedative hypnotics	
Drug-Inferred Health State – Opioids and Prenatal Vitamins and Medications Used in Pregnancy	Enhanced point-of-sale alert for concurrent use of opioids and prenatal vitamins and for concurrent use of opioids and medications used in pregnancy (e.g., doxylamine/pyridoxine)	



Concurrent Drug Utilization Review (cDUR) programs

 This custom message does not require the pharmacist to enter appropriate NCPDP codes to receive an approved claim

Retrospective Drug Utilization Review (rDUR) programs

The rDUR program analyzes claims on a daily basis and sends communications to prescribers.

Abused Medications DUR Program

- Daily identification of members who are getting multiple opioid prescriptions from multiple prescribers and/or filling at multiple pharmacies
- Also identifies members with chronic early refill attempts, overlapping LAOs, overlapping SAOs, high daily doses of opioids, a diagnosis of opioid overdose with an opioid fill, overlapping opioid and MAT medications, overlapping opioid and a benzodiazepine, overlapping opioid and an antipsychotic medication, overlapping opioid, muscle relaxant and benzodiazepine and overlapping opioid and opioid potentiator
- Patient-specific information sent to all prescribers with medication fill history for the last 4 months

Pharmacy Lock-In Program

- Pharmacy lock-in programs vary by state, however, all include filling of opioids by multiple prescribers at multiple pharmacies as an inclusion criterion for the program
- Members chosen for the program will be locked into 1 pharmacy for all of their medications for a period of 1 year. Lock-in periods vary by state.

Utilization Management (UM) programs

UM programs promote appropriate use, help reduce costs and ultimately, help improve the health status of members.

Cumulative 90 Morphine Milligram Equivalent (MME) Limit

- Point-of-sale dosage limit for all opioid products up to 90 MME
- Prevents cumulative opioid doses above the preset threshold from processing
- Prior authorization required for doses above the preset threshold

LAO Prior Authorization

Prior authorization requires:

- Attestation of appropriate use and monitoring
- Step through short-acting opioid (non-cancer pain) and step through preferred LAOs
- If appropriate, step through neuropathic pain alternatives (non-cancer pain)

New-to-Therapy SAO Edit

- Point-of-sale limits for members who are opioid naïve (no opioid claims in the last 60 days)
- Point-of-sale limits include maximum of 30 MME and/or 20 units per fill on select SAOs
- Prior authorization required to exceed these quantities for opioidnaïve members

Cough and Cold Products Containing Opioid Components

- Quantity per fill of 120 mL (units) is applied as well as a 30-day maximum quantity of 360 mL (units)
- Prior authorization is required for those under age 18

Transmucosal Fentanyl Product Prior Authorization

 Prior authorization requires documentation of pain due to cancer and patient is already receiving opioids



Utilization Management (UM) programs

UM programs promote appropriate use, help reduce costs and ultimately, help improve the health status of members.

Buprenorphine MAT	 No prior authorization required for preferred products with	
Prior Authorization	appropriate diagnosis at point-of-sale	
Overdose Prevention (naloxone)	 No prior authorization is required for preferred naloxone products (generic naloxone injection, Narcan[®] Nasal Spray) 	

Evidence-Based Prescribing programs

Focuses on outreach to prescribers identified as outliers

Fraud/Waste/Abuse Evaluation	Retrospective controlled substance claims analysisIdentifies outlier opioid prescribers		
Miscellaneous			
Substance Use Disorder Helpline	 24/7 helpline: 855-780-5955. For members or caregivers staffed by licensed behavioral health providers Reference: liveandworkwell.com 		
Miscellaneous – Drug Enforcement Agency (DEA) License Edit	 Verifies DEA is active and matches scheduled medication in the claim 		
Miscellaneous – Refill-Too-Soon Threshold	 Increases the refill-too-soon threshold to 90% on opioids and other controlled substances CII-V 		

Abbr	eviations		
APAP	Acetaminophen	MME	Morphine Milligram Equivalent
CDC	Centers for Disease Control and Prevention	PA	Prior Authorization
cDUR	Concurrent Drug Utilization Review	rDUR	Retrospective Drug Utilization Review
DEA	Drug Enforcement Agency	SAOs	Short-Acting Opioids
LAOs	Long-Acting Opioids	UM	Utilization Management
MAT	Medication-Assisted Treatment		



We're here to help

For more information, please call Provider Services at 888-362-3368.

How to submit prior authorizations

- Online: Use the Prior Authorization and Notification tool in Link. For more information, go to UHCprovider.com/paan.
- Phone: Call 800-310-6826
- Fax: Fax your completed form to 866-940-7328
- Pharmacy Prior Authorization forms are available at UHCprovider.com > Menu > Health Plans by State choose your state > Medicaid (Community Plan) > Pharmacy Resources and Physician-Administered Drugs > Pharmacy Prior Authorization

