

Ilumya® (Tildrakizumab-Asmn)

Policy Number: CS2023D0074L
Effective Date: May 1, 2023

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Applicable Codes	2
Background	3
Clinical Evidence	3
U.S. Food and Drug Administration	3
References	3
Policy History/Revision Information	4
Instructions for Use	4

Commercial Policy
• Ilumya® (Tildrakizumab-Asmn)

Application

This Medical Benefit Drug Policy does not apply to the states listed below; refer to the state-specific policy/guideline, if noted:

State	Policy/Guideline
Indiana	Immunomodulators for Inflammatory Conditions (for Indiana Only)
Kansas	Refer to the state’s Medicaid clinical policy
Louisiana	Refer to the state’s Medicaid clinical policy
North Carolina	None
Ohio	Ilumya® (Tildrakizumab-Asmn) (for Ohio Only)
Pennsylvania	Refer to the state’s Medicaid clinical policy
Washington	Refer to the state’s Medicaid clinical policy

Coverage Rationale

Ilumya (tildrakizumab), to be used as a self-administered, subcutaneous injection for the treatment of plaque psoriasis, should be obtained under the pharmacy benefit.

Initial Therapy

Ilumya (tildrakizumab) is proven and medically necessary for provider administration for the treatment of moderate to severe plaque psoriasis when the following criteria are met:

- Submission of medical records (e.g., chart notes, laboratory values) documenting all of the following:
 - Diagnosis of chronic moderate to severe plaque psoriasis; and
 - Greater than or equal to 3% body surface area involvement, palmoplantar, facial, genital involvement, or severe scalp psoriasis^{1,2,3,6,8}; and
 - One of the following:
 - Both of the following:
 - History of failure, contraindication, or intolerance to one of the following topical therapies:⁴

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Anthralin
- Coal tar

and

- History of failure to a 3 - month trial of methotrexate at maximally indicated dose, unless contraindicated or clinically significant adverse effects are experienced^{6,7}

or

- Patient has been previously treated with a biologic or targeted synthetic DMARD FDA-approved for the treatment of plaque psoriasis [e.g., Cimzia (certolizumab), Humira (adalimumab), Otezla (apremilast), Skyrizi (risankizumab - rzaa), Stelara (ustekinumab), Tremfya (guselkumab)].

and

- History of failure, contraindication, or intolerance to three biologic DMARDs FDA-approved for the treatment of plaque psoriasis (document drug, date, and duration of trial); and
- Physician attestation that the patient is unable to self-administer or there is no competent caregiver to administer the drug. Physician must submit explanation; and
- Patient is not receiving Ilumya in combination with any of the following:¹
 - Biologic DMARD [e.g., Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab), Cosentyx (secukinumab), Orencia (abatacept)]
 - Janus kinase inhibitor [e.g., Xeljanz (tofacitinib)]
 - Phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)]

and

- Dosing is in accordance with the United States Food and Drug Administration approved labeling; and
- Prescribed by or in consultation with a dermatologist; and
- Initial authorization will be for no longer than 12 months

Continuation Therapy

Ilumya (tildrakizumab) will be reauthorized for provider administration based on all of the following criteria:

- Documentation of positive clinical response to Ilumya therapy; and
- Physician attestation that the patient is unable to self-administer or there is no competent caregiver to administer the drug. Physician must submit explanation; and
- Patient is not receiving Ilumya in combination with any of the following:
 - Biologic DMARD [e.g., Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab)]
 - Janus kinase inhibitor [e.g., Xeljanz (tofacitinib)]
 - Phosphodiesterase 4 (PDE4) inhibitor [e.g., Otezla (apremilast)]
- and
- Dosing is in accordance with the United States Food and Drug Administration approved labeling; and
- Reauthorization will be for no longer than 12 months

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
J3245	Injection, tildrakizumab, 1 mg

Diagnosis Code	Description
L40.0	Psoriasis vulgaris

Background

Ilumya (tildrakizumab) is a humanized IgG1/k monoclonal antibody that selectively binds to the p19 subunit of IL - 23 and inhibits its interaction with the IL - 23 receptor. IL-23 is a naturally occurring cytokine that is involved in inflammatory and immune responses. Tildrakizumab inhibits the release of pro-inflammatory cytokines and chemokines.

Clinical Evidence

Plaque Psoriasis

Ilumya (tildrakizumab) is an interleukin - 23 antagonist indicated for the treatment of adults with moderate to severe plaque psoriasis who are candidates for systemic therapy or phototherapy.^{1,9}

Professional Societies

American Academy of Dermatology (AAD)

In 2019, the AAD and the National Psoriasis Foundation published updated treatment guidelines for the management and treatment of psoriasis with biologic therapies. In regards to tildrakizumab and/or IL - 23 inhibitors, the guidelines state:

- Tildrakizumab is recommended as a monotherapy treatment option in adult patients with moderate-to-severe plaque psoriasis.
- The recommended dose is 100 mg given by in office physician-administered subcutaneous injection at week 0 and week 4 and every 12 weeks thereafter.
- There is no evidence to support combination of tildrakizumab with topical or systemic therapies, but there is no reason to consider such combination unsafe.
- Definitive response (positive or negative) to treatment with IL - 23 antagonists is best ascertained after 12 weeks of continuous therapy. Consider dose escalation in partially responding patients. Consider the addition of other modalities (such as topical corticosteroids or vitamin D analogues, methotrexate, or ultraviolet B light) in partially responding patients. Although there are no published data supporting combination therapy for the IL - 23 inhibitors, there is no reason to consider such combination therapy unsafe.
- The effect of guselkumab on solid tumor or lymphoreticular malignancy, when used as monotherapy for moderate-to-severe psoriasis, is unknown. Large long-term follow-up studies are necessary to more fully define the risk of cancer associated with IL - 23 inhibitors.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Ilumya (tildrakizumab) is an interleukin-23 antagonist indicated for the treatment of adults with moderate to severe plaque psoriasis who are candidates for systemic therapy or phototherapy.

References

1. Ilumya [prescribing information]. Cranbury, NJ: Sun Pharmaceutical Industries, Inc.; July 2020.
2. Menter A, Gottlieb A, Feldman SR, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. J Am Acad Dermatol 2008; 58(5):826-50.
3. Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Psoriatic arthritis: Overview and guidelines of care for treatment with an emphasis on the biologics. J Am Acad Dermatol 2008;58(5):851-64.

4. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 3. Guidelines of care for the management and treatment of psoriasis with topical therapies. J Am Acad Dermatol 2009;60(4):643-59.
5. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Guidelines of care for the treatment of psoriasis with phototherapy and photochemotherapy. J Am Acad Dermatol 2010;62(1):114-35.
6. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. J Am Acad Dermatol 2009;61(3):451-85.
7. Nast A, et al; European S3-Guidelines on the systemic treatment of psoriasis vulgaris – update 2015 – short version – EFF in cooperation with EADV and IPC, J Eur Acad Derm Venereol 2015;29:2277-94.
8. Menter A, Korman NJ, Elmets CA, Feldman SR, Gelfand JM, Gordon KB, Guidelines of care for the management of psoriasis and psoriatic arthritis: section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011 Jul;65(1):137-74.
9. Reich K, Papp KA, Blauvelt A, et al. Tildrakizumab versus placebo or etanercept for chronic plaque psoriasis (reSURFACE 1 and reSURFACE 2): results from two randomized controlled, phase 3 trials. Lancet. 2017 Jul 15;390(10091):276-288.
10. Reich K, Warren RB, Iversen L, et al. Long-term efficacy and safety of tildrakizumab for moderate-to-severe psoriasis: pooled analyses of two randomized phase III clinical trials (reSURFACE 1 and reSURFACE 2) through 148 weeks. Br J Dermatol. 2019 Jun 19.
11. Elewski B, Menter A, Crowley J, et al. Sustained and Continuously Improved Efficacy of Tildrakizumab in Patients with Moderate-to-Severe Plaque Psoriasis. J Dermatolog Treat. 2019 Jul 3:1-19.
12. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. J Am Acad Dermatol. 2019 Apr;80(4):1029-1072.

Policy History/Revision Information

Date	Summary of Changes
05/01/2023	<p>Coverage Rationale</p> <ul style="list-style-type: none"> ● Revised coverage criteria for initial therapy: <ul style="list-style-type: none"> ○ Replaced criterion requiring “history of failure, contraindication, or intolerance to <i>two of the following preferred biologic products: Humira (adalimumab), Enbrel (etanercept), and/or Cimzia (certolizumab)</i>” with “history of failure, contraindication, or intolerance to <i>three biologic DMARDs FDA-approved for the treatment of plaque psoriasis (document drug, date, and duration of trial)</i>” ○ Removed criterion requiring one of the following: <ul style="list-style-type: none"> ▪ History of 6 - month trial of Cosentyx (secukinumab) with moderate clinical response yet residual disease activity ▪ Both of the following: <ul style="list-style-type: none"> – History of intolerance or adverse event to Cosentyx – Physician attests that in their clinical opinion the same intolerance or adverse event would not be expected to occur with Ilumya <p>Supporting Information</p> <ul style="list-style-type: none"> ● Archived previous policy version CS2022D0074K

Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check

the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.